

Cochrane Database of Systematic Reviews

Vitamin D for the management of asthma (Review)

Martineau AR, Cates CJ, Urashima M, Jensen M, Griffiths AP, Nurmatov U, Sheikh A, Griffiths CJ
Martineau AR, Cates CJ, Urashima M, Jensen M, Griffiths AP, Nurmatov U, Sheikh A, Griffiths CJ. Vitamin D for the management of asthma. <i>Cochrane Database of Systematic Reviews</i> 2016, Issue 9. Art. No.: CD011511. DOI: 10.1002/14651858.CD011511.pub2.

www.cochranelibrary.com



[Intervention Review]

Vitamin D for the management of asthma

Adrian R Martineau¹, Christopher J Cates², Mitsuyoshi Urashima³, Megan Jensen⁴, Alex P Griffiths⁵, Ulugbek Nurmatov⁶, Aziz Sheikh⁷, Chris J Griffiths¹

¹Asthma UK Centre for Applied Research, Barts and The London School of Medicine and Dentistry, Queen Mary University of London, London, UK. ²Population Health Research Institute, St George's, University of London, London, UK. ³Division of Molecular Epidemiology, Jikei University School of Medicine, Tokyo, Japan. ⁴Centre for Asthma and Respiratory Diseases, University of Newcastle, Newcastle, Australia. ⁵Centre for Primary Care and Public Health and Asthma UK Centre for Applied Research, Barts and The London School of Medicine and Dentistry, Queen Mary University of London, London, UK. ⁶Division of Population Medicine, School of Medicine, the National Centre for Population Health and Wellbeing Research, Cardiff University, Cardiff, UK. ⁷Asthma UK Centre for Applied Research, Usher Institute of Population Health Sciences and Informatics, The University of Edinburgh, Edinburgh, UK

Contact address: Adrian R Martineau, Asthma UK Centre for Applied Research, Barts and The London School of Medicine and Dentistry, Queen Mary University of London, London, UK. a.martineau@qmul.ac.uk.

Editorial group: Cochrane Airways Group.

Publication status and date: Edited (no change to conclusions), published in Issue 4, 2019.

Citation: Martineau AR, Cates CJ, Urashima M, Jensen M, Griffiths AP, Nurmatov U, Sheikh A, Griffiths CJ. Vitamin D for the management of asthma. *Cochrane Database of Systematic Reviews* 2016, Issue 9. Art. No.: CD011511. DOI: 10.1002/14651858.CD011511.pub2.

Copyright © 2019 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

ABSTRACT

Background

Several clinical trials of vitamin D to prevent asthma exacerbation and improve asthma control have been conducted in children and adults, but a meta-analysis restricted to double-blind, randomised, placebo-controlled trials of this intervention is lacking.

Objectives

To evaluate the efficacy of administration of vitamin D and its hydroxylated metabolites in reducing the risk of severe asthma exacerbations (defined as those requiring treatment with systemic corticosteroids) and improving asthma symptom control.

Search methods

We searched the Cochrane Airways Group Trial Register and reference lists of articles. We contacted the authors of studies in order to identify additional trials. Date of last search: January 2016.

Selection criteria

Double-blind, randomised, placebo-controlled trials of vitamin D in children and adults with asthma evaluating exacerbation risk or asthma symptom control or both.

Data collection and analysis

Two review authors independently applied study inclusion criteria, extracted the data, and assessed risk of bias. We obtained missing data from the authors where possible. We reported results with 95% confidence intervals (CIs).

Main results

We included seven trials involving a total of 435 children and two trials involving a total of 658 adults in the primary analysis. Of these, one trial involving 22 children and two trials involving 658 adults contributed to the analysis of the rate of exacerbations requiring systemic corticosteroids. Duration of trials ranged from four to 12 months, and the majority of participants had mild to moderate asthma. Administration of vitamin D reduced the rate of exacerbations requiring systemic corticosteroids (rate ratio 0.64, 95% CI 0.46 to 0.90;



680 participants; 3 studies; high-quality evidence), and decreased the risk of having at least one exacerbation requiring an emergency department visit or hospitalisation or both (odds ratio (OR) 0.39, 95% CI 0.19 to 0.78; number needed to treat for an additional beneficial outcome, 27; 963 participants; 7 studies; high-quality evidence). There was no effect of vitamin D on % predicted forced expiratory volume in one second (mean difference (MD) 0.48, 95% CI -0.93 to 1.89; 387 participants; 4 studies; high-quality evidence) or Asthma Control Test scores (MD -0.08, 95% CI -0.70 to 0.54; 713 participants; 3 studies; high-quality evidence). Administration of vitamin D did not influence the risk of serious adverse events (OR 1.01, 95% CI 0.54 to 1.89; 879 participants; 5 studies; moderate-quality evidence). One trial comparing low-dose versus high-dose vitamin D reported two episodes of hypercalciuria, one in each study arm. No other study reported any adverse event potentially attributable to administration of vitamin D. No participant in any included trial suffered a fatal asthma exacerbation. We did not perform a subgroup analysis to determine whether the effect of vitamin D on risk of severe exacerbation was modified by baseline vitamin D status, due to unavailability of suitably disaggregated data. We assessed two trials as being at high risk of bias in at least one domain; neither trial contributed data to the analysis of the outcomes reported above.

Authors' conclusions

Whilst we are confident that Vitamin D reduced the risk of asthma exacerbation in these trials (high quality GRADE assessment), we recognise that there is uncertainty about how these findings might be applied in practice. More research is needed to clarify whether there is a difference in effect between adults and children and with respect to asthma severity, baseline vitamin D status and doses.

PLAIN LANGUAGE SUMMARY

Vitamin D to prevent asthma attacks

Review question

Does vitamin D prevent asthma attacks or improve control of asthma symptoms or both?

Background

Low blood levels of vitamin D (the 'sunshine vitamin') have been linked to an increased risk of asthma attacks in children and adults with asthma. Several clinical trials have been conducted to test whether vitamin D might prevent asthma attacks and improve control of asthma symptoms in children and adults, but results from studies with the most scientifically sound designs have not previously been evaluated as a group.

Included studies

We included seven trials involving 435 children and two trials involving 658 adults in the review from searches run up to January 2016. Of these, one trial involving 22 children and two trials involving 658 adults contributed to the analysis of the rate of severe asthma attacks. Study duration ranged from four to 12 months, and the majority of those taking part had mild or moderate asthma. All of the studies compared vitamin D with placebo.

Key results

People given vitamin D experienced fewer asthma attacks needing treatment with oral steroids. The average number of attacks per person per year went down from 0.44 to 0.28 with vitamin D (high-quality evidence). Vitamin D reduced the risk of attending hospital with an acute asthma attack from 6 per 100 to around 3 per 100 (high-quality evidence).

Vitamin D had little or no effect on lung function or day-to-day asthma symptoms (high-quality evidence). We found that vitamin D did not increase the risk of serious adverse events at the doses that were tested (moderate-quality evidence).

We based all of these findings on studies judged to be of high quality.

Conclusion

Vitamin D has been found to offer some protection against severe asthma attacks in adults with mild to moderate asthma. Further trials focusing on children and people who experience frequent severe asthma attacks are needed before definitive clinical recommendations can be made.